



Dr. Charles Gardner, Medical Officer of Health Dr. Colin Lee, Associate Medical Officer of Health Dr. Lisa Simon, Associate Medical Officer of Health

Universal Influenza Immunization Program (UIIP) 2023-2024

Attention: Long-Term Care Facilities, Retirement Homes, Hospital Pharmacies, Hospital Occupational,

Health Nurses, Infection Control Practitioners, Ontario Health Central

Date: September 15, 2023

Information about the local rollout of COVID-19 XBB vaccine and Respiratory Syncytial Virus (RSV) vaccine will be shared via separate Public Health Alerts as soon as we receive more information from the Ministry of Health. As per the Ministry of Health press release this week, RSV vaccine will be available for residents of long-term care homes, elder care lodges and some retirement home residents this fall. To support facilities planning, it is expected that RSV vaccine will need to be given at least 14 days following COVID or flu vaccine, so administration is likely to be later in October.

The following Influenza vaccines will be available publicly funded in Ontario this flu season:

	QIV		QIV-HD	TIV-adj
Age Group	FluLaval Tetra	Fluzone® Quadrivalent	Fluzone® High-Dose Quadrivlalent	Fluad®
6 months to 64 years	✓	✓		
≥ 65 years	✓	✓	✓	✓

Strains of influenza in the vaccine products offered in the 2023/2024 UIIP:

Influenza Strains	Egg-based QIVs	Egg-based TIVs (Fluad®)
A/Victoria/4897/2022 (H1N1)pdm09-like virus;	NEW For 2023/2024	NEW For 2023/2024
A/Darwin/9/2021 (H3N2)-like virus;	✓	✓
B/Austria/1359417/2021 (B/Victoria lineage)-like virus;	~	~
B/Phuket/3073/2013 (B/Yamagata lineage)-like virus.	~	



Who Should Receive the Influenza Vaccine?

Per NACI recommendations for influenza and to optimize co-administration with COVID-19 vaccine, health care workers, first responders, and the following individuals at high risk of influenza-related complications or who are more likely to require hospitalization, should be prioritized to receive the influenza vaccine as soon as vaccine is available:

- Residents and staff of congregate living settings (e.g. chronic care facilities, retirement homes)
- People ≥ 65 years of age
- All pregnant individuals
- All children 6 months to 4 years of age
- Individuals who are from a First Nation, Inuit or Métis community, and/or who self-identify as First Nation, Inuit
 or Métis, and their household members
- Members of racialized and other equity deserving communities
- Individuals 6 months of age and older with the following underlying health conditions:
 - o Cardiac or pulmonary disorders
 - Diabetes mellitus or other metabolic disease
 - o Cancer
 - Conditions or medication which compromise the immune system
 - o Renal disease
 - o Anemia or hemoglobinopathy
 - o Neurologic or neurodevelopment conditions
 - Morbid obesity (body mass index of ≥40)
 - Children and adolescents (6 months to 18 years) undergoing treatment with acetylsalicylic acid for long periods

The influenza vaccine is recommended for all people six months of age and older without contraindications, however, per NACI, individuals in the following three groups are particularly recommended to receive the influenza vaccine:

- Individuals capable of transmitting influenza to those listed above and/or to infants under 6 months of age:
 - o Health care workers and other care providers in facilities and community settings
 - Household contacts (adults and children) of individuals at high risk of influenza related complications
 - o Persons who provide care to children ≤ 4 years of age
 - o Members of a household expecting a newborn during the influenza season
 - o Those who provide services within a closed or relatively closed setting to persons at high risk of influenza related complications (e.g. crew on a ship)
- People who provide essential community services
- Poultry industry workers

How Many Doses of Influenza Vaccine are Needed for Protection?

Age		Number of doses recommended	
6 months to under 9	Not previously immunized with any influenza vaccine in their lifetime	2 doses at least 4 weeks apart ¹	
years of age	Previously immunized with at least one dose of any influenza vaccine in their lifetime	1 dose	
9 years of age	and older	1 dose	

¹ It is not necessary to use the same vaccine product for both doses.

Publicly Funded Influenza Vaccines for Anyone 6 Months of Age and Older

	Quadrivalent Inactivated Vaccines		
Vaccine product	FluLaval Tetra	Fluzone® Quadrivalent	
Age indication	≥ 6 months	≥ 6 months	
Manufacturer	GSK	Sanofi Pasteur	
Vaccine Type	Egg-based	Egg-based	
Micrograms of hemagglutinin	15 µg	15 µg	
Dosage	0.5 mL	0.5 mL	
Format	MDV (multi-dose vial – stable for 28 days once punctured)	once punctured)	
		PFS (pre-filled syringe)	
Route	Intramuscular (IM) injection	Intramuscular (IM) injection	
Most common allergens ¹	Egg protein ²	Egg protein ²	
	Thimerosal (multi-dose vial only)	Thimerosal (multi-dose vial only)	

¹ Any component in a vaccine may be a potential allergen. This table identifies the most common allergens.

Which Influenza Vaccine Should Individuals ≥ 65 Years of Age Receive?

The QIV, QIV-HD and TIV-adj <u>all protect against the flu</u>, and the most important thing is for older adults to be vaccinated as soon as vaccine is available. **Do not delay vaccination to wait for a particular product.**

For individual-level decision making, NACI states that when available, high-dose (QIV-HD) should be used over standard-dose (QIV) inactivated influenza vaccine, given the burden of influenza A (H3N2) disease and the good evidence of better protection compared to standard-dose in adults 65 years of age and older.

However, NACI states that in the absence of a specific product, any of the available age-appropriate influenza vaccines should be used. **There is NO PREFERENTIAL RECOMMENDATION** for the use of QIV-HD versus TIV-adj vaccine for this age group. To date, no studies have directly compared QIV-HD and TIV-adj formulations or TIV-adj and QIV formulations. The following information should be considered when discussing vaccine options:

- There is insufficient evidence to make a preferential recommendation between QIV-HD and TIV-adj. There is good evidence of QIV-HD providing better protection compared to QIV standard dose.
- In considering use of TIV-adj and QIV standard dose, given the increased burden of disease associated with influenza A(H3N2) in older adults, better protection against influenza A(H3N2) afforded by the TIV-adj may be more important, especially in those with multiple co-morbid conditions and compromised health status. However, most importantly, do not delay vaccination to wait for a particular product.

² The National Advisory Committee on Immunization (NACI) indicates that egg allergy is not a contraindication for influenza vaccination and that that egg-allergic individuals may be vaccinated against influenza using the full dose of any age-appropriate product. See section IV of the Canadian Immunization Guide chapter on Influenza and statement on seasonal influenza vaccine for 2018-2019 for studies supporting the NACI recommendation for egg-allergic individuals (www.phac-aspc.gc.ca/naci-ccni/#rec).

Publicly Funded Influenza Vaccines for Anyone 65 Years of Age and Older Only

	High-Dose Quadrivalent Inactivated Vaccine	Adjuvanted Trivalent Inactivated Vaccine		
Vaccine product	Fluzone® High-Dose Quadrivalent	Fluad [®]		
Age indication	≥ 65 years	≥ 65 years		
Manufacturer	Sanofi Pasteur	Seqirus		
Vaccine Type	Egg-based	Egg-based		
Micrograms of hemagglutinin	60 µg	15 µg		
Dosage	0.7 mL	0.5 mL		
Adjuvant	No	Yes		
Format	PFS (pre-filled syringe)	PFS (pre-filled syringe)		
Route	Intramuscular (IM) injection	Intramuscular (IM) injection		
Most common allergens ¹	• Egg protein ²	 Egg protein² Neomycin Kanamycin 		

¹ Any component in a vaccine may be a potential allergen. This table identifies the most common allergens.

Fluzone® Quadrivalent and Fluzone® High-Dose Quadrivalent (QIV-HD) are different products. Fluzone® High-Dose Quadrivalent (QIV-HD) is only authorized for those 65 years of age and older. Please use caution when administering Fluzone® products to ensure that the right vaccine is being administered to the right person.

Co-administration with Other Vaccines

Influenza vaccines may be given on the same day, or at any time before or after other vaccines (including COVID-19 vaccines) for anyone 6 months of age and older.

No studies have been conducted that have assessed the co-administration of **Shingrix** with adjuvanted or high-dose influenza vaccines. With Fluad®, it is unknown how the adjuvants may interact when Shingrix is co-administered.

Reporting Adverse Events

An <u>Adverse Event Following Immunization (AEFI) form</u> must be filled out and faxed to the Immunization Program's fax line at **705-726-3962** for follow-up as soon as possible following any unexpected adverse events. See Public Health Ontario's AEFI Reporting Fact Sheet for more information.

² The National Advisory Committee on Immunization (NACI) indicates that egg allergy is not a contraindication for influenza vaccination and that that egg-allergic individuals may be vaccinated against influenza using the full dose of any age-appropriate product. See section IV of the Canadian Immunization Guide chapter on Influenza and statement on seasonal influenza vaccine for 2018-2019 for studies supporting the NACI recommendation for egg-allergic individuals (www.phac-aspc.gc.ca/naci-ccni/#rec).

Vaccine Ordering Information

Health care providers are only to immunize those in the high-risk groups outlined on page 2 until the annual influenza program is officially launched by the Ministry of Health, which usually occurs ~November 1st. Once the Ministry of Health launches the annual UIIP, vaccine can be administered to the general population.

Facilities	Order Due By	Ready for Pick up	Product	Maximum Quantity
Hospitals For high-risk patients (per list on pg. 2) & staff	Wed Sept 27 th by 3 p.m.	Wed Oct 4 th at 8:30 a.m.	QIV-HD QIV *See below	Based on assessment of your facilities needs
LTCHs & RHomes For residents, staff and essential caregivers	Wed Sept 27 th by 3 p.m.	Wed Oct 4 th at 8:30 a.m.	QIV-HD QIV **See below	Based on bed numbers
Reorders	Wed Oct 11 th by 3 p.m.	Wed Oct 18 th at 8:30 a.m.	QIV-HD QIV	Based on local supply

Hospitals*

Order as needed for patients considered high-risk as per list on page 2 and staff. Order QIV-HD for any patients who are 65+, and order QIV for the remaining.

Pre-filled syringes (PFS) will be shipped, so please only order what your fridge can safely store. If you would prefer multi-dose vials please note this on your order form. PFS do not come with needle tips, so ensure you have some on hand to administer.

Long Term Care Homes and Retirement Homes**

Order the number of doses of QIV-HD for residents who are 65+ and order QIV for any residents under 65. Vaccine packaging is large, so please only order what your fridge can safely store, and re-order as needed.

If you have a medical directive in place to immunize staff and essential caregivers, order the required number of QIV doses required. QIV for staff may be either pre-filled syringe or multi-dose vial based on SMDHU inventory availability.

QIV-HD will be provided in a pre-filled syringe but does not come with a needle tip so please ensure you have enough needle tips in stock.

Only influenza vaccine orders submitted on this year's order form will be filled. The 2023-2024 Influenza Vaccine Order Form is available on the <u>SMDHU Health Professional portal</u>. Be sure to choose the appropriate form based on which health unit office you pick up vaccine from. Any orders submitted prior to today on last year's order forms will be disregarded.

Each time you order influenza vaccine, you must include your temperature log sheet(s) for the previous four weeks. Temperature readings must be done and recorded in the logbook twice a day, each day your facility is open (including dates and times of readings). These temperature log sheets are reviewed by SMDHU prior to vaccine being released.

If there is a reason you are ordering a large amount of vaccine (i.e., you have a clinic booked), please note this on the order form so that SMDHU staff can take this into account when reviewing your order. **Supply for large clinics will not be available until November**.

Ensure your vaccine cooler and your vaccine fridge will be able to accommodate any influenza vaccine order placed. You may need to make more than 1 trip to pick up your order or bring an additional hard-sided cooler and thermometer as we do not have any extra supplies to loan out.

Additional Resources

For more information, please visit our website at www.smdhu.org/hpportal or contact the Immunization Program at 705-721-7520 or toll free at 1-877-721-7520

- For questions regarding vaccine orders: ext. 8808 or vaccineorders@smdhu.org
- For all other HCP questions: ext. 8806 or hc.vpd@smdhu.org